

Basic Information

Product Name	Anti-HSP70 Antibody (Clone#OTI2C12)		
Gene Name	HSPA1A		
Source	Mouse		
Clonality	Monoclonal		
Isotype	IgG1		
Species Reactivity	human, mouse, rat, monkey		
Tested Application	WB, IHC, ICC/IF, FCM		
Contents	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.		
Immunogen	Full length human recombinant protein of human HSPA1A (NP_005336) produced in HEK293T cell.		
Concentration	500 ug/ml		
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)		
Observed MW	70 kDa		
Dilution Ratios	Western blot (WB):	1:2000	
	Immunohistochemistry (IHC):	1:50	
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:100	
	Flow cytometry (FCM):	1:100	

Storage

Stable for 12 months from date of receipt. Store at -20°C as received.


Background Information

This intronless gene encodes a 70kDa heat shock protein which is a member of the heat shock protein 70 family. In conjunction with other heat shock proteins, this protein stabilizes existing proteins against aggregation and mediates the folding of newly translated proteins in the cytosol and in organelles. It is also involved in the ubiquitin-proteasome pathway through interaction with the AU-rich element RNA-binding protein 1. The gene is located in the major histocompatibility complex class III region, in a cluster with two closely related genes which encode similar proteins.

Reference

Anti-HSP70 Antibody (Clone#OTI2C12)被引用在1文献中。

Selected Validation Data



Western blot analysis of anti-HSP70 antibody (MA00949). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,

Lane 2: human Hela whole cell lysates,

Lane 3: human A431 whole cell lysates,

Lane 4: human MCF-7 whole cell lysates,

Lane 5: rat kidney tissue lysates,

Lane 6: rat C6 whole cell lysates,


Lane 7: mouse kidney whole cell lysates,

Lane 8: mouse NIH/3T3 whole cell lysates.


After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with mouse anti-HSP70 antigen affinity purified monoclonal antibody (MA00949) and probed with a goat anti-mouse IgG-HRP secondary antibody (Catalog # BA1050).


The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for HSP70 at approximately 70 kDa. The expected band size for HSP70 is at 70 kDa.



Immunohistochemical staining of paraffin-embedded liver tissue within the normal limits using anti-HSP70 mouse monoclonal antibody (MA00949). (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, MA00949, Dilution 1:50)



Immunofluorescent staining of HepG2 cells using anti-HSP70 mouse monoclonal antibody (MA00949) (1:50).



Flow cytometric analysis of permeabilized HeLa cells, using anti-HSP70 antibody, compared to an isotype control (green), and a PBS control (blue) (1:100).