#### **Product datasheet**

### **Anti-FOSL1 Antibody (Clone#OTI12F9)**

Catalog Number: MA03927



Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

Basic Information	
Product Name	Anti-FOSL1 Antibody (Clone#OTI12F9)
Gene Name	FOSL1
Source	Mouse
Clonality	Monoclonal
Isotype	lgG2b
Species Reactivity	human, mouse, rat
Tested Application	IHC, WB
Contents	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Immunogen	Full-length protein expressed in 293T cell transfected with human FOSL1 expression vector
Concentration	500 ug/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Observed MW	29.4 kDa
Dilution Ratios	Western blot (WB): 1:2000 Immunohistochemistry (IHC):1:50

# **Storage**

Stable for 12 months from date of receipt. Store at -20°C as received.

# **Background Information**

The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation.

## Reference

Anti-FOSL1 Antibody (Clone#OTI12F9)被引用在1文献中。

### Anti-FOSL1 Antibody (Clone#OTI12F9)


Catalog Number: MA03927




Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

# **Selected Validation Data**



Immunohistochemical staining of paraffin-embedded prostate tissue within the normal limits using anti-FOSL1mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, MA03927, Dilution 1:50)



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FOSL1 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FOSL1 (Cat#MA03927).