

Basic Information

Product Name	Anti-IP10/CXCL10 Antibody
Gene Name	CXCL10
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human
Tested Application	WB, ELISA
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.
Immunogen	E. coli-derived human IP10 recombinant protein (Position: V22-P98). Human IP10 shares 70.1% and 72.7% amino acid (aa) sequence identity with mouse and rat IP10, respectively.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	11 kDa
Dilution Ratios	Western blot (WB): 1:500-2000 Enzyme linked immunosorbent assay (ELISA):1:100-1000


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

C-X-C motif chemokine 10 (CXCL10), also known as Interferon gamma-induced protein 10 kDa (IP-10) or small-inducible cytokine B10, is a protein that in humans is encoded by the CXCL10 gene. It is a small cytokine belonging to the CXC chemokine family. The gene for CXCL10 is located on human chromosome 4 in a cluster among several other CXC chemokines. Luster et al. (1987) reported the isolation of an interferon-inducible gene that encodes a 98-amino acid protein called IP10. The chemokine IFN-gamma-inducible protein 10 (IP-10; CXCL10), a CXC chemokine, and its receptor, CXCR3, were found to be overexpressed in lymph node cells of EAMG rats. IP10 inhibits bone marrow colony formation, has antitumor activity in vivo, is a chemoattractant for human monocytes and T cells, and promotes T cell adhesion to endothelial cells.

Selected Validation Data



Western blot analysis of IP10/CXCL10 using anti-IP10/CXCL10 antibody (PB0751). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: HELA whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-IP10/CXCL10 antigen affinity purified polyclonal antibody (PB0751) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for IP10/CXCL10 at approximately 11 kDa. The expected band size for IP10/CXCL10 is at 11 kDa.