# Product datasheet Anti-FSHR Antibody Catalog Number: PB1120



Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

<b>Basic Information</b>	
Product Name	Anti-FSHR Antibody
Gene Name	FSHR
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	E. coli-derived human FSH Receptor recombinant protein (Position: C18-N187). Human FSH Receptor shares 91.2% and 90% amino acid (aa) sequence identity with mouse and rat FSH Receptor, respectively.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	78 kDa
Dilution Ratios	Western blot (WB):1:500-2000

#### **Storage**

12 months from date of receipt, -20°C as supplied.

# **Background Information**

The follicle-stimulating hormone receptor or FSH receptor (FSHR) is a transmembrane receptor that interacts with the follicle-stimulating hormone (FSH) and represents a G protein-coupled receptor (GPCR). This FSHR gene is mapped to chromosome 2p21 by fluorescence in situ hybridization. The protein encoded by this gene belongs to family 1 of G-protein coupled receptors. It is the receptor for follicle stimulating hormone and functions in gonad development. Mutations in this gene cause ovarian dysgenesis type 1, and also ovarian hyperstimulation syndrome. Alternative splicing results in multiple transcript variants.

# Reference

Anti-FSHR Antibody被引用在5文献中。




**BOSTER BIOLOGICAL TECHNOLOGY** 

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator, East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com

# **Selected Validation Data**



Western blot analysis of FSHR using anti-FSHR antibody (PB1120). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat testis tissue lysates,

Lane 2: rat ovary tissue lysates,

Lane 3: mouse testis tissue lysates,

Lane 4: mouse ovary tissue lysates,

Lane 5: human Hela whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-FSHR antigen affinity purified polyclonal antibody (PB1120) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for FSHR at approximately 78 kDa.

The expected band size for FSHR is at 78 kDa.