

Basic Information

| | |
|---------------------------|---|
| Product Name | Anti-GPX1 Antibody |
| Gene Name | GPX1 |
| Source | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Species Reactivity | human, mouse, rat |
| Tested Application | WB, IHC |
| Contents | 500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol. |
| Immunogen | A synthetic peptide corresponding to a sequence in the middle region of human GPX1, different from the related mouse sequence by six amino acids and from the related rat sequence by five amino acids. |
| Concentration | 500 ug/ml |
| Purification | Immunogen affinity purified. |
| Observed MW | 22 kDa |
| Dilution Ratios | Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user. |

Storage

12 months from date of receipt, -20°C as supplied.

Background Information


Glutathione peroxidase 1, also known as, GPX-1 is an enzyme that in humans is encoded by the GPX1 gene. It is mapped to 3p21.31. This gene encodes a member of the glutathione peroxidase family, consisting of eight known glutathione peroxidases (Gpx1-8) in humans. Glutathione peroxidase functions in the detoxification of hydrogen peroxide, and is one of the most important antioxidant enzymes in humans. It has been reported that the protein encoded by this gene protects from CD95-induced apoptosis in cultured breast cancer cells and inhibits 5-lipoxygenase in blood cells, and its overexpression delays endothelial cell growth and increases resistance to toxic challenges. GPX1

is one of only a few proteins known in higher vertebrates to contain selenocysteine, which occurs at the active site of glutathione peroxidase and is coded by the nonsense (stop) codon TGA.

Reference

Anti-GPX1 Antibody被引用在4文献中。

Selected Validation Data



Western blot analysis of GPX1 using anti-GPX1 antibody (PB9203). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human THP-1 whole cell lysates,

Lane 2: rat lung tissue lysates,

Lane 3: rat liver tissue lysates,

Lane 4: mouse lung tissue lysates,

Lane 5: mouse liver tissue lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-GPX1 antigen affinity purified polyclonal antibody (PB9203) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for GPX1 at approximately 22 kDa. The expected band size for GPX1 is at 22 kDa.

Product datasheet

Anti-GPX1 Antibody

Catalog Number: **PB9203**


BOSTER[®]

antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com



IHC analysis of GPX1 using anti-GPX1 antibody (PB9203) .

GPX1 was detected in a paraffin-embedded section of human colorectal adenocarcinoma tissue. The tissue section was incubated with rabbit anti-GPX1 Antibody (PB9203) at a dilution of 1:200 and developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB (Catalog # AR1027) as the chromogen.